

08 616371

MAR 15 1996
DB/CAE/ach
3/15/96METHODS FOR PRODUCING AND USING S-NITROSOHEMOGLOBINSBACKGROUND OF THE INVENTION

Interactions of hemoglobin (Hb) with small ~~diffuseable~~ ^{diffusible} ligands, such as O₂, CO₂ and NO, are known to occur at its metal centers and amino termini. The O₂/CO₂ delivery functionalities, which arise in the lung and systemic microvasculature, are allosterically controlled. Such responsiveness to the environment is not known to apply in the case of NO. Specifically, it is thought that Hb(Fe) is involved in limiting NO's sphere of action (Lancaster, J.R., *proc. Natl. Acad. Sci. USA*, 91:8137-8141 (1994); Wood and Garthwaite, *J. Neuropharmacol.*, 33:1235-1244 (1994)), but that NO does not modify the functional properties of Hb to any physiologically significant degree. Kinetic modeling based on this assumption, however, predicts that the vast majority of free NO in the vasculature should be scavenged by Hb (Lancaster 1994). Accordingly, the steady-state level of NO may fall below the K_m for target enzymes such as guanylate cyclase (Lancaster 1994), if not in the unperturbed organism, then with oxidant stress such as that found in atherosclerosis. These considerations raise the fundamental question of how NO exerts its biological activity.

One answer to this paradox may be found in the propensity of nitric oxide to form S-nitrosothiols (RSNOs) (Gaston, B. et al., *Proc. Natl. Acad. Sci. USA*, 90:10957-10961 (1993)), which retain NO-like vasorelaxant activity (Stamler, J.S., et al., *Proc. Natl. Acad. Sci. USA*, 89:444-448 (1992)), but which are not subject to the diffusional constraints imposed by the high concentration of Hb in the blood. In particular, the NO group of RSNOs possesses nitrosonium (NO⁺) character that distinguishes it from NO itself. Indeed, it is increasingly appreciated that RSNO's have the capacity to elicit certain functions that NO is

a incapable of (DeGroote, M.A. et al., *Proc. Natl. Acad. Sci. USA*, 92:6399-6403 (1995); Stamler, J.S., *Cell*, 78:931-936 (1994)). Moreover, consideration has been given to the possibility that -SNO groups in proteins may serve a

5 signaling function, perhaps analagous to phosphorylation (Stamler, J.S. et al., *Proc. Natl. Acad. Sci. USA*, 89:444-448 (1992); Stamler, J.S. *Cell*, 78:931-926 (1994)).

a Although S-nitrosylation of proteins can regulate protein function (Stamler, J.S. et al., *Proc. Natl. Acad. Sci. USA*, 10 89:444-448 (1992); Stamler, J.S., *Cell*, 78:931-936 (1994)), the identification of S-nitrosoproteins within cells -- the *sine qua non* of a regulatory posttranslational modification -- has heretofore not been demonstrated.

a Hemoglobin is a tetramer comprised of two alpha and 15 two beta subunits. In human Hb, each subunit contains one heme, while the beta (β) subunits also contain highly reactive -SH groups (cys β 93) (Olson, J.S., *Meth. in Enzym.*, 76:631-651 (1981); Antonini & Brunori, *In Hemoglobin and Myoglobin in Their Reactions with Ligands*, American 20 Elsevier Publishing Co., Inc., New York, pp. 29-31 (1971)). These cysteine residues are highly conserved among species although their function has remained elusive.

NO (nitric oxide) is a biological "messenger molecule" which decreases blood pressure and inhibits platelet 25 function, among other functions. NO freely diffuses from endothelium to vascular smooth muscle and platelet and across neuronal synapses to evoke biological responses. Under some conditions, reactions of NO with other components present in cells and in serum can generate toxic 30 intermediates and products at local concentrations in tissues which are effective at inhibiting the growth of infectious organisms. Thus, it can be seen that a method of administering an effective concentration of NO or biologically active forms thereof would be beneficial in 35 certain medical disorders.

SUMMARY OF THE INVENTION

The invention relates to a method of forming SNO-Hb by reaction of Hb with S-nitrosothiol in a procedure which avoids oxidation of the heme. The invention also relates to a method of therapy for a condition in which it is desired to oxygenate, to scavenge free radicals, or to release NO⁺ groups to tissues. SNO-Hb in its various forms and combinations thereof (oxy, deoxy, met; S-nitrosylated to various extents) can be administered to a mammal in this method. Thiols can also be administered to enhance the transfer of NO groups. Examples of conditions to be treated by SNO-Hb therapy include ischaemic injury, hypertension, angina, reperfusion injury and inflammation.

BRIEF DESCRIPTIONS OF THE DRAWINGS

a 15 Figures 1A-1D are spectrographs of different forms of Hb as described in Example 1.

Figure 2A is a graph showing formation, with time, of SNO-Hb by S-nitrosylation.

20 Figure 2B is a graph showing the decomposition, with time, of oxy and deoxy forms of SNO-Hb.

Figure 3A is a graph showing the loading of red blood cells with S-nitrosocysteine, over time. The inset is a series of spectrographs of forms of Hb as described in Example 3.

25 Figure 3B is a series of tracings recording isometric tone of a rabbit aortic ring following treatment of the aortic ring with various agents as described in Example 3.

Figure 4A is a graph of change in tension of a rabbit aortic ring versus concentration of the Hb used in the experiment.

30 Figure 4B is a graph of change in tension of a rabbit aortic ring versus concentration of the Hb used in the experiment, where glutathione was also added to test the effect as compared to Figure 4A.

Figure 4C is a graph of the ratio of S-nitrosogluthathione formed/starting SNO-Hb concentration versus time, showing rates of NO group transfer from oxy and met forms of Hb to glutathione.

5 Figure 4D is a graph of S-nitrosothiols exported from loaded red blood cells over time.

Figure 5 is a graph showing the mean arterial blood pressure in rats after they received various doses of oxyHb (▲), SNO-oxyHb (■), or SNO-metHb (●).

10 Figures 6A-6F are a series of tracings recording blood pressure (Figures 6A and 6B), coronary artery diameter (Figures 6C and 6D) and coronary artery flow (Figures 6E and 6F), after administration of S-nitrosohemoglobin to anesthetized dogs.

15 DETAILED DESCRIPTION OF THE INVENTION

Roles for Hemoglobin in Physiology

The increase in SNO-Hb content of red cells across the pulmonary circuit (right ventricular inport-left ventricle) suggests that the Hb molecule is S-nitrosylated in the
20 lung. Selective transfer of the NO group from endogenous RSNOs found in lung (Gaston, et al. (1993)) and blood (Scharfstein, et al. (1994)) to SH groups of Hb, substantiate these findings. Nonetheless, the mechanism(s) of S-nitrosylation operative in vivo is not known. The
25 corresponding decline in Hb(FeII)NO levels across the pulmonary bed reveals a role for the lung either in the elimination of NO or in its intramolecular transfer from heme to cys β 93. Taken in aggregate, these data extend the list of function-regulating interactions of Hb with small
30 molecules within the respiratory system, previously known to include the elimination of CO and CO₂ and uptake of O₂. Since oxygenation of Hb leads to structural changes that increase the NO-related reactivity of cys β 93, O₂ may be regarded as an allosteric effector of Hb S-nitrosylation.

This is a newly discovered allosteric function for the protein.

The arterial-venous difference in SNO-Hb concentration suggests that the protein acts as an NO group donor in the systemic circulation. Indeed, there is good indication that SNO-Hb functions in regulation of vasomotor tone. In the microcirculation, where control of blood pressure is achieved, erythrocytes come in intimate contact with endothelial surfaces. Under these conditions, Hb can predispose the vasculature to increases in arterial resistance by sharply decreasing the steady state level of free NO (Lancaster, J.R., (1994)). This rationale is believed to contribute to the increases in blood pressure that occur with infusion of cell-free Hbs (Vogel, W.M., et al. (1986); Olsen, S.B., et al. (1996)). The transient nature of such hypertensive responses, however, would be consistent with the subsequent formation of SNO-Hb which counteracts this effect, evidenced by its lowering of blood pressure at naturally occurring concentrations. Thus, the capacity of the erythrocyte to support the synthesis and metabolism of SNO-Hb may well be important for normal blood flow.

It is implicit from this line of reasoning that mammals must have adopted unique molecular mechanisms to ensure adequate NO delivery in the microcirculation. Our results suggest that Hb may have evolved both electronic and conformational switching mechanisms to achieve NO homeostasis. Specifically, NO scavenging by the metal center(s) of SNO-Hb(FeII)O₂ would be sensed through its conversion to met (FeIII) (Figure 1B). This electronic event would effect decomposition of SNO-Hb with NO group release (Figures 3A, 3B, 4A). In this manner, the NO-related activity of SNO-Hb would be partly determined by the amount of NO scavenged. Changes in O₂ tension might also function to regulate NO delivery, as we observed that

NO release was facilitated by deoxygenation. It is intriguing to speculate that this allosteric effect may operate to limit the tissue O₂ deficit. In this scenario, NO group release would serve to regulate capillary blood flow to enhance O₂ delivery.

S-nitrosothiol groups in proteins have been implicated in NO metabolism and in regulation of cellular functions (Stamler, J.S., et al., *Proc. Natl. Acad. Sci USA*, 89:444-448 (1992); Stamler, J.S., *Cell*, 78:931-936 (1994)). The identification of SNO-Hb in erythrocytes is the first demonstration of an intracellular S-nitrosoprotein and gives further credence to their role in cellular regulation. The question arises as to how SNO-Hb relaxes blood vessels when any free NO released would be scavenged instantaneously by Hb itself (Lancaster, J.R., (1994)). Noteworthy in this regard are studies showing that RSNO activity involves nitrosyl (NO⁺) transfer to thiol acceptors (Scharfstein, J.S., et al., (1994); Arnette and Stamler, (1995); Stamler, J.S., et al., *Proc. Natl. Acad. Sci. USA*, 89:7674-7677 (1992)), which serve to protect the NO-related activity from inactivation at metal centers. Findings of this study indicate that S-nitrosothiol/thiol exchange with glutathione (forming GSNO) is likely to occur within erythrocytes, and to be influenced by the oxidation state of heme and its occupation by ligand. ^{DeGroote} ~~DeGroote~~ and coworkers have shown that certain activities of GSNO in bacteria require transport of intact dipeptide (i.e S-nitrosocysteinyglycine) across the cell membrane (DeGroote, M.A., et al., (1995)). The data presented here expand this paradigm to include eukaryotic cells. GSNO, or related thiol carriers exported by erythrocytes (Kondo and Beutler, *Methods ⁱⁿ Enzymology*, Packer, L., ed., Academic Press, 252:72-83, (1995)), might also initiate ^{signaling} ~~signalling~~ in or at the ^{plasma membrane} ~~plasmalemma~~ (Stamler, J.S., *Cell*, 78:931-936 (1994)), given reports of thiol-dependent activation of

potassium channels by EDRF (Bolotina, V.M., et al., *Nature*, 368:850-853 (1994)). Alternative possibilities also merit consideration. In particular, reports that Hb associates with red cell membranes via cys β 93 (Salhany and Gaines, *Trends in Biochem. Sci.*, Jan 13-15 (1981)). This would place Hb in position to donate the NO group directly to contacting endothelial surfaces, perhaps via SNO/SH exchange. Indeed, cell surface interactions appear to be operative in signaling mediated by other S-nitrosoproteins (Stamler, J.S., et al., *Proc. Natl. Acad. Sci. USA*, 89:444-448 (1992); Stamler, J.S., *Cell*, 78:931-936 (1994)).

The highly conserved cys β 93 residues in Hb have been shown to influence the oxygenation and propensity for oxidation of the metal center, and to affect the physicochemical behavior of the protein (Garel, C., et al., (1982); Jocelyn, P.C., (1972); Craescu, C.T., et al., (1986); Mansouri, A., *Biochem. Biophys. Res. Commun.*, 89:441-447 (1979)). Nonetheless, their long sought after physiological function has remained a mystery. Our studies suggest new sensory and regulatory roles for Hb, in which cys β 93 functions in transducing NO-related signals to the vessel wall. In particular, the physiological function of cys β 93, which is invariant in all mammals and birds, is to deliver under allosteric control, NO-related biological activity that cannot be scavenged by heme. Thus, these data bring to light a dynamic circuit for the NO group in which intraerythrocytic Hb participates as both a sink and a donor, depending on the particular microenvironment of the circuit. Such observations may provide answers to paradoxes that arise from conceptual frameworks based solely on diffusional spread and reaction of free NO (Lancaster, J.R., (1994); Wood and Garthwaite, (1994)); and may have implications that extend to other thiol- and metal-containing (heme) proteins, such as nitric oxide synthase and guanylate cyclase.

Finally, the discoveries reported here may have direct therapeutic implications. Specifically, concerns over loss of NO-related activity due to inactivation by blood Hb (Lancaster, J.R., (1994)) are obviated by the presence of an RSNO subject to allosteric control. SNO-Hb is free of the adverse hypertensive properties of cell-free Hb preparations that result from NO scavenging at the metal centers. A cell-free Hb solution that ^{mimics} ~~mimicks~~ blood by containing SNO-Hb can be used as a blood substitute.

10 Further embodiments

The subject invention relates to a method of loading cells with a nitrosating agent as exemplified for red blood cells as in Figure 3A, but which can be accomplished in more general ways. Suitable conditions for pH and for the temperature of incubation are, for example, a range of pH 7-9, with pH 8 being preferred, and a temperature range of 25 to 37°C. For red blood cells, short incubation times of 1 to 3 minutes are preferred for limiting the formation of S-nitrosylated forms of Hb. However, intracellular concentrations of 1 mM nitrosating agent can be reached.

The nitrosating agent should be a good donor of NO⁺ and ~~should be able to~~ diffuse through the cell membrane of the target cell type. That is, it must be of low molecular weight, in contrast to S-nitroso proteins. Examples are S-nitroso-N-acetylcysteine, S-nitrosocysteinylglycine, S-nitrosocysteine, S-nitrosohomocysteine, organic nitrates and nitrites, metal nitrosyl complexes and other related nitrosating agents as defined in "Donors of Nitrogen Oxides" pp. 71-119 In Methods in Nitric Oxide Research (Feelisch, M. and Stamler, J.S., eds.) Wiley, Chichester, U.K. (1996), the contents of which chapter are hereby incorporated by reference in their entirety. Nitrosating agents have differential activities for different reactive groups on metal-containing proteins. A nitrosating agent

can be chosen for minimal oxidation of the heme iron of Hb, and maximum activity in nitosylating thiol groups such as found on cysteine.

Such low molecular weight nitrosating agents can be used in red blood cells to deliver NO-related activity to tissues. Treatment of red blood cells with nitrosating agent further serves to increase the O₂ delivery capacity of red blood cells. Such treatment of red blood cells also allows for the scavenging of oxygen free radicals throughout the circulation. Therefore, it is possible to load red blood cells with S-nitrosothiol, for example, by a process outside a patient's body after removal of whole blood (as a minimal method of isolating the red blood cells) and then to reintroduce the red blood cells into the same patient), thereby allowing the treatment of a number of types of diseases and medical disorders, such as those which are characterized by abnormal O₂ metabolism of tissues, oxygen-related toxicity, abnormal vascular tone, abnormal red blood cell adhesion, or abnormal O₂ delivery by red blood cells. Such diseases can include, but are not limited, to ischaemic injury, hypertension, shock, angina, stroke, reperfusion injury, acute lung injury, sickle cell anemia, schistosomiasis and malaria. The use of such "loaded" red blood cells also extends to blood substitute therapy and the preservation of living organs, as organs for transplantation, for example. In some cases, it may be appropriate to treat a patient with loaded red blood cells originating from a different person.

A particular illustration of the mechanism of the treatment method is presented here by considering sickle cell anemia. Sickle cell patients suffer from frequent vascular occlusive crises which manifest in clinical syndromes such as the acute chest syndrome and hepatic dysfunction. Both endothelial cell dysfunction, resulting in a clotting diathesis as well as dysfunction intrinsic to

the red blood cell, are central to disease pathogenesis. At the molecular level, the increased expression of vascular adhesion molecules such as VCAM promote the adhesion of sickled red blood cells, containing abnormal hemoglobin. It follows that decreasing cytokine expression on endothelial cells, promoting endothelial function and attenuating red cell sickling, are key therapeutic objectives. However, currently used therapies have been generally unsuccessful.

10 In this novel method for loading red blood cells with intracellular NO-donor S-nitrosothiols, the effect is to increase oxygen affinity -- which in and of itself should attenuate red blood cell sickling -- and to endow the red blood cell with vasodilator and antiplatelet activity, 15 which should reverse the vasoocclusive crisis. Moreover, nitric oxide should attenuate the expression of adhesion molecules on endothelial cell surfaces, thus restoring endothelial function.

Herein is described a novel therapeutic approach to 20 the treatment of sickle cell disease which involves loading of red blood cells with S-nitrosothiols or other nitrosating agents. Two examples of therapeutic approaches are given. In the first, the patient's own red blood cells would be S-nitrosylated extracorporeally (yielding "loaded" 25 red blood cells) and then given to the patient. The second approach is to directly administer to a patient an agent such as S-nitrosocysteine, which is permeable to red blood cells.

For some diseases or disorders, the administration of 30 NO-loaded red blood cells is especially desirable. Upon a change from the oxygenated to the deoxygenated state, or upon a change in the oxidation state of the heme Fe from the reduced state (FeII) to the oxidized (FeIII) state, NO is released from the thiol groups of hemoglobin, and is 35 rapidly transferred to glutathione to form S-

nitrosoglutathione. Red blood cells are known to have a high concentration of glutathione. S-nitrosoglutathione efficiently delivers NO to tissues.

In another aspect, the invention is a method for the treatment of infection by administering to an infected mammal an agent which causes S-nitrosylation of thiol groups within the cells which are the target of such agent. For example, an S-nitrosothiol to which lymphocytes are highly permeable can be administered to a patient infected with HIV. Such treatment for HIV can also be used excorporeally, to blood isolated from the patient. In another application, the infection is bacterial, and the S-nitrosothiol to be used as an anti-bacterial agent is one to which the target bacterial cells are highly permeable, as compared to the permeability properties of the host cells. (See, for example De Groote, M.A., et al., *Proc. Natl. Acad. Sci. USA* 92:6399-6403 (1995).) Alternatively, nitrosothiols can be used to treat *Plasmodium falciparum* within red blood cells.

Another embodiment of the invention is a method for specifically modifying a protein containing one or more metal atoms so that the protein becomes S-nitrosylated at one or more thiol groups without modifying the metal, as by changing the oxidation state or causing the metal atoms to bind NO. This can be accomplished by the use of a reagent which possesses NO⁺ character, such as a nitrosothiol (see, for instance, Example 4A), which reacts specifically with thiol groups of a protein in which metal is bound.

For hemoglobin, the nitrosation method does not affect the heme. SNO-Hb (SNO-Hb(FeII)O₂) can be synthesized from Hb(FeII)O₂ with up to 2 SNO groups per tetramer without oxidation of the heme Fe from FeII to FeIII. In contrast, when Hb(FeII)O₂ is incubated with excess nitric oxide or nitrite, methemoglobin (HbFe[III]) forms rapidly (Example 1B) and to a significant extent. When Hb[FeII] is

incubated with nitric oxide, NO binds rapidly to the heme, forming Hb(FeII)NO to a significant extent (Example 1A).

Although rates of formation of SNO-Hb(FeII)O_2 from Hb(FeII)O_2 are more rapid (see Example 2A), the
5 corresponding SNO-deoxyHb form can also be made by incubation of S-nitrosoglutathione or S-nitrosocysteine, for example, with Hb(FeII) , yielding SNO-Hb(FeII) , as in Example 1C.

The effects of the various forms of Hb on vasodilation
10 -- constriction, dilation or a neutral effect -- depend on three factors: whether 1) the Fe of the heme is oxidized, 2) O_2 is bound at the heme (that is, the oxygenation state, dictated by the conformation of the protein as R state or T state), and 3) thiol is present in sufficient concentration
15 to facilitate the transfer of NO^+ .

The importance of the first factor is shown in Figure 4A. Hb(FeII)O_2 and SNO-Hb[FeII]O_2 act as vasoconstrictors, but SNO-Hb[FeIII] (met form) where FeII has been oxidized to FeIII) acts as a vasodilator. Figure 4A shows that SNO-Hb[FeII]O_2 with oxygen bound at the heme, and with a ratio
20 of $\text{SNO/Hb}=2$, acts as a powerful vasoconstrictor.

SNO-Hb(FeII) is also a vasodilator. Figure 2B illustrates the second factor in demonstrating that rates of RSNO decomposition and transfer are much faster for SNO-Hb in the deoxy state than for SNO-Hb in the oxy state.
25

It can be seen how the NO^+ -donating properties of SNO-Hb depend on oxygen concentrations. SNO-Hb releases oxygen at sites of low oxygen concentration or under oxidizing conditions. SNO-Hb releases its NO group(s) to cause
30 vasodilation either due to 1) oxidation of the heme Fe to FeIII or 2) loss of the O_2 from the heme by deoxygenation. It is shown in Figure 2B that NO is transferred off SNO-Hb best in the deoxy state. In ischaemia, SNO-Hb deoxygenates, rapidly followed by the loss of NO. It can
35 be seen from the data that SNO-metHb having a ratio of 1

SNO/SNO-metHb is a more powerful vasodilator than SNO-oxyHb having a ratio of 2 SNO/SNO-oxyHb. It should be noted that S-nitrosylation of Hb induces the R state (oxy conformation). Thus, it follows that 1 SNO-oxyHb molecule
5 having a ratio of 1 SNO/SNO-oxyHb is less potent than 10 molecules of SNO-oxyHb having a ratio of 0.1 SNO/SNO-oxyHb.

The third factor is illustrated by the results shown in Figure 4B. These results demonstrate potentiation by thiol of the vasodilator effect of SNO-Hb(FeII)O₂ and SNO-
10 Hb(FeIII). Transfer of NO⁺ from SNO-Hb to low molecular weight nitrosothiols is more efficient when Hb is in the deoxy state compared to the oxy state (Figure 2B) or in the met state compared to the oxy state (Figure 4C).

NO is released or transferred as NO⁺ (nitrosyl cation)
15 from SNO-Hb. The SNO groups of SNO-Hb have NO⁺ character. Transfer of NO⁺ from SNO-Hb occurs most efficiently to small thiols, such as glutathione, and is most efficient when the heme is oxidized (SNO-metHb) or the SNO-Hb is in the deoxy state.

20 One embodiment of the invention resulting from these findings is a method of therapy that enhances the transfer of NO⁺ from SNO-Hb to small thiols, thereby delivering NO biological activity to tissues, by the coadministration of small molecular weight thiols, along with a form of SNO-Hb,
25 to a mammal in need of the physiological effects of NO. To further increase the effect of NO release it is preferred that the SNO- forms of metHb or deoxyHb (or an equivalent conformation or spin state) be administered with the thiol (See Figure 2B, for example.) A mixture of SNO-metHb and
30 SNO-oxyHb, and possibly also thiol, can also be used. The composition and proportion of these components depends on the disease state. For example, to achieve both enhanced O₂ delivery and NO delivery, SNO-oxyHb can be used. Where no further delivery of O₂ is desirable, as in acute
35 respiratory distress syndrome, for example, the SNO- forms

of methHb and deoxyHb are especially preferred. Alternatively, the ratios of SNO/Hb can be regulated to control O₂ release.

The vessel ring bioassay data of Figure 4A agree well with the *in vivo* data of Figure 5. The results of the experiments described in Example 5 confirm that Hb(FeII)O₂ (oxyHb) causes an increase in blood pressure *in vivo*, as it did also *in vitro*. SNO-Hb(FeIII) (SNO-metHb) causes a decrease in blood pressure *in vivo* as well as *in vitro*. SNO-Hb(FeII)O₂ (SNO-oxyHb) has a negligible effect on blood pressure *in vivo* in contrast to the increase in tension seen in the corresponding vessel ring bioassay. For SNO-oxyHb the *in vivo* effect is neutral. This may be explained by the constrictive effect caused by NO becoming bound to the heme being compensated by the release of NO upon deoxygenation. Therefore, SNO-oxyHb can deliver O₂ with minimal effect on blood pressure.

With knowledge of the results herein it is possible to synthesize Hb proteins with predicted NO-releasing properties which will constrict, dilate, or have no effect on blood vessels. An additional option is the choice between making oxygenated or deoxygenated forms to administer for medical conditions in which O₂ delivery is desirable, or undesirable, respectively.

It is possible to produce a variety of modified Hbs having specific desired properties of O₂ and NO delivery. For example, Hb in the R state (oxyHb) can be converted to the T state (deoxyHb) by a number of known methods. This can be done, for example, by reaction of Hb with inositol hexaphosphate. It is also known to those skilled in the art that Hb in the R state can be made, for example, by treating Hb with carboxypeptidase. Similarly, it is known that methHb can be synthesized using ferricyanide or nitrite.

Producing Hb molecules which are locked in the T state allows the synthesis of RSNO-Hb which remains in a form that is a biologically active donor of NO, rather than a carrier of NO. Hb which is locked in the R state can be
5 used as a substrate for the synthesis of RSNO-Hb which carries a maximum amount of NO per molecule.

Another embodiment of the invention is a blood substitute comprising one or more forms of Hb which have been specifically S-nitrosylated to some extent at one or
10 more thiol groups of the Hb, in order to regulate O₂ release and NO release. Conditions to be treated include those in which NO or O₂ delivery is desired, those in which NO or O₂ utilization is desired, or those in which NO or O₂ is in excess. For example, in a medical condition which is
15 characterized by the presence of an excess of oxygen free radicals and excess NO[•], both the heme of SNO-Hb and NO released by SNO-Hb serve to trap oxygen free radicals. The heme Fe is oxidized in the process of scavenging oxygen free radicals and NO[•], and NO is released from the oxidized
20 Hb by donation to a thiol, in the form of RSNO⁺ (nontoxic). Inflammation and reperfusion injury, for example, are characterized by excess NO production and an excess of oxygen free radicals. Forms of Hb scavenge oxygen radicals and free NO, converting NO to forms that are not toxic.

25 A further embodiment of the invention is a method of therapy for a condition that would benefit from the delivery of NO biological activity or O₂ or both, based on the administration of a blood substitute comprising a form of SNO-Hb. For example, SNO-Hb is useful to treat
30 myocardial infarction. SNO-Hb has the effect of donating NO, keeping blood vessels open. SNO-Hb deoxygenates at low oxygen tension, delivering oxygen and releasing NO at the same site, thereby causing vasodilation. (See Example 7 and Figures 6A-6F.) These effects can be augmented by
35 also administering thiol, either simultaneously with SNO-

Hb, or before or after. For the purpose of treating myocardial infarction, for example, a high concentration or dose of SNO-Hb that has a low ratio of SNO/SNO-Hb is appropriate. Alternatively, SNO-metHb can be used for this purpose.

In another aspect, the invention is a method of enhancing NO-donor therapy by coadministering SNO-Hb together with a nitroso-vasodilator (nitroglycerin, for example) which would be otherwise consumed by the conversion of oxyHb to metHb in Hb which has not been S-nitrosylated.

The term hemoglobin or Hb as used herein includes variant forms such as mutant forms, chemically modified forms, genetically altered forms, such as fusion proteins, and truncated forms. It also includes Hbs of all species and variant forms thereof. The biological and/or chemical properties of these variant Hbs may be different from those of hemoglobins which are found naturally occurring in animals.

It will be appreciated that NO exists in biological systems not only as nitric oxide gas, but also in various redox forms and as biologically active adducts of nitric oxide such as S-nitrosothiols, which can include S-nitrosoproteins, S-nitroso-amino acids and other S-nitrosothiols (Stamler, J.S. Cell 78:931-936 (1994)).

A blood substitute can be a biologically compatible liquid which performs one or more functions of naturally occurring blood found in a mammal such as oxygen carrying and/or delivery, NO carrying and/or delivery, and the scavenging of free radicals. A blood substitute can also comprise one or more components of such a liquid which, when infused into a mammal, perform one or more functions of naturally occurring blood. Examples of blood substitutes include preparations of various forms of hemoglobin. Such preparations may also include other

biologically active components, such as a low molecular weight thiol or nitrosothiol.

The compounds and therapeutic preparations of this invention to be used in medical treatment are intended to be used in therapeutic amounts, in suitable compositions, which can be determined by one of skill in the art. Modes of administration are those known in the art which are most suitable to the affected site or system of the medical disorder.

a 10 Suitable pharmaceutical carriers can be combined ~~with~~ with active ingredients employed in a therapeutic preparation, if necessary. It will be appreciated that the actual amounts of the active components in a specific case will vary according to the specific component being
15 utilized, the particular composition formulated, the mode of administration and the age, weight and condition of the patient, for example. Dosages for a particular patient can be determined by one of ordinary skill in the art using conventional considerations, (e.g. by means of an
20 appropriate, conventional pharmacological protocol).

The present invention is more specifically illustrated in the following examples, which are not intended to be limiting in any way.

EXAMPLES

25 Example 1: Interactions of NO and RSNO with Hb

a It was observed that naturally occurring N-oxides, such as NO and RSNOs (Gaston, B., *et al.* (1993; Scharfstein, J.S., *et al.*, *J. Clin. Invest.*, 94:1432-1439 (1994); Clancy, R.M., *et al.*, *Proc. Natl. Acad. Sci. USA*,
30 91:3680-3684 (1994)), differed markedly in their reactions with Hb. NO bound very rapidly to deoxyHb (Hb[FeII]),

forming relatively stable $\text{Hb}[\text{FeII}]\text{NO}$ complexes (Figure 1A), and converted oxyHb ($\text{Hb}[\text{FeII}]\text{O}_2$) to methemoglobin ($\text{Hb}[\text{FeIII}]$) and nitrate (Figure 1B), confirming previous reports (Olson, (1981); Toothill, C., *Brit. J. Anaesthy.*, 5 39:405-412 (1967)). In contrast, RSNOs were found to participate in transnitrosation reactions with sulfhydryl groups of Hb, forming S-nitrosohemoglobin (SNO-Hb), and did not react with the heme centers of either deoxyHb or $\text{Hb}(\text{FeII})\text{O}_2$ (Figures 1C and 1D).

10 A. Interaction of NO with deoxyHb

Conversion of deoxyHb ($\text{Hb}[\text{FeII}]$) to $\text{Hb}(\text{FeII})\text{NO}$ is observed upon incubation of $\text{Hb}(\text{FeII})$ with increasing concentrations of nitric oxide. a. Deoxy Hb. b, c, d. Reaction mixtures of NO and $\text{Hb}(\text{FeII})$ in ratios of 1:1, 2:1 and 10:1, respectively. The reaction product $\text{Hb}(\text{FeII})\text{NO}$ formed essentially instantaneously on addition of NO (i.e. within instrument dead time).

B. Interaction of NO with oxyHb

Conversion of oxyHb ($\text{Hb}[\text{Fe}[\text{II}]\text{O}_2$) to metHb ($\text{HbFe}[\text{III}]$) is observed upon incubation of oxyHb with increasing concentrations of NO. a. oxy Hb. b, c, d. Reaction mixtures containing NO and oxyHb in ratios of 1:1, 2:1 and 10:1, respectively. Methemoglobin formation occurred instantaneously on addition of NO (i.e. within instrument 25 dead time).

C. Interaction of S-nitrosothiols with deoxyHb

Conversion of $\text{Hb}(\text{FeII})$ to SNO-Hb(FeII) is observed upon incubation of either GSNO (shown) or S-nitrosocysteine (CYSNO) with deoxy Hb. There is little (if any) 30 interaction of RSNO with the heme functionalities of Hb. a. deoxyHb. b, c, d. Reaction mixtures of GSNO and $\text{Hb}(\text{FeII})$ in ratios of 1:1, 2:1 and 10:1, respectively. Spectra were

taken after 60 min of incubation in *b*, *c*, and 15 min in *d*. Further analysis of reaction products revealed the formation of moderate amounts of SNO-Hb in all cases. Yields of SNO-Hb (S-NO/Hb) in *b*, *c*, and *d* at 60 min were
5 2.5%, 5% and 18.5%, respectively. (See Figure 1D and Figure 2A.)

D. Interaction of S-nitrosothiols with oxyHb

Conversion of Hb(FeII)O₂ to SNO-Hb(FeII)O₂ is observed upon incubation of either GSNO (shown) or CYSNO with oxyHb.
10 There is little (if any) reaction of GSNO (or CYSNO) at the heme centers of Hb(FeII)O₂. Specifically, the capacity for O₂ binding to heme is unaffected by RSNOs. *a*. oxyHb. *b*, *c*, *d*. Reaction mixtures of GSNO and oxyHb in ratios of 1:1, 2:1, and 10:1, respectively. Spectra were taken after 60
15 min of incubation in the spectrophotometer. Further analysis of reaction products revealed the formation of SNO-Hb in all cases. Yields of SNO-Hb in spectra *b*, *c* and *d* were 5%, 10% and 50% (S-NO/Hb), respectively. In 5 other determinations, the yield of S-NO/Hb was 0.37 ± 0.06
20 using GSNO (pH 7.4, 10-fold excess over Hb) and ~2 SNO/tetramer (1.97 ± 0.06) using CYSNO (*vida infra*). These last data are in agreement with reports that human HbA contains 2 titratable SH groups.

Methods

a
25 Human HbA₀ was purified from red cells as previously described (Kilbourn, R.G., et al., *Biochem. Biophys. Res. Comm.*, 199:155-162 (1994)). Nitric oxide solutions were rigorously degassed and purified according to standard procedure (Beckman, J. S., et al., *Methods in Nitric Oxide*
30 *Research*, Feelisch and Stamler, eds., Wiley Chichester, U.K. (1996)) and saturated solutions were transferred in air tight syringes. Deoxygenation of Hb was achieved by addition of excess dithionite (NO studies) or by reduction

- of Hb(FeII)O₂ through evacuation in Thunberg tubes (RSNO studies; as RSNOs react with dithionite). RSNOs were synthesized as previously described (Gaston, B., et al., (1993); Arnelle and Stamler, (1995)) Incubations with HbA₀ were made in phosphate buffer, pH 7.4, 0.5 mM EDTA. Quantifications of SNO-Hb were made according to the method of Saville (Gaston, B., et al., (1993); Stamler, J.S., et al., *Proc. Natl. Acad. Sci. USA*, 90:444-448 (1992)) after purification of protein with Sephadex G-25 columns. The Saville method, which assays free NO_x in solution, involves a diazotization reaction with sulfanilamide and subsequent coupling with the chromophore N-(naphthyl)ethylenediamine. No low molecular weight S-NO complexes survived this purification and all activity was protein precipitable. The reactions and spectra were carried out using a Perkin Elmer UV/Vis Spectrometer, Lambda 2S.

Example 2: Allosteric Function of O₂ in Regulation of Hb S-nitrosylation

- Oxygenation of Hb is associated with conformational changes that increase the reactivity of cysβ93 to alkylating reagents (Garel, C., et al., *J. Biochem.*, 123:513-519 (1982); Jocelyn, P.C., *Biochemistry of the SH Group*, Academic Press, London, pp. 243 (1972); Craescu, C.T., et al., *J. Biol. Chem.*, 261:14710-14716 (1986)). The physiological importance of this effect was never established. We observed that rates of S-nitrosylation of Hb were markedly dependent on conformational state. In the the oxy conformation (R state), S-nitrosylation was more rapid than in the deoxy conformation (T state) (Figure 2A). The rate of S-nitrosylation was accelerated in both conformations by alkaline conditions (i.e. rate at pH 9.2 > pH 7.4), which would tend to expose the cysβ93 that is otherwise screened from reaction by the C-terminal histidine 146β. The salt bridge (asp β94 --- his β146)

tying down the histidine residue is loosened at high pH. These data suggest that the increase in thiol reactivity associated with the R state derives, at least in part, from improved NO access rather than a conformation-induced
5 change in pK.

a A. Oxygenation accelerates S-nitrosylation of Hb

Rates of Hb S-nitrosylation by S-nitrosocysteine (CYSNO) are faster in the oxy conformation (Hb[FeII]O₂) than in the deoxy state (Hb[FeII]).

10 Methods

a Incubations were performed using 10-fold excess CYSNO over protein (50 μM) in aerated 2% borate, 0.5 mM EDTA (oxyHb), or in a tonometer after rapid O₂ evacuation (deoxyHb). At shown times, samples were rapidly desalted
15 ^{across} ~~across~~ G-25 columns (preequilibrated with phosphate buffered saline, 0.5 mM EDTA, pH 7.4) to remove CYSNO, and analyzed for SNO-Hb by the method of Saville (Stamler, J.S., et al., *Proc. Natl. Acad. Sci. USA*, 89:444-448 (1992)).

20 B. Deoxygenation accelerates denitrosylation of Hb

Rates of RSNO decomposition (and transfer) are much faster in the deoxy conformation [SNO-Hb(FeII)] than in the oxy state [SNO-Hb(FeII)O₂]. The decomposition of SNO-Hb(FeII) is further accelerated by the presence of excess
25 glutathione. Within the dead time of our measurements (~15 sec) a major fraction of SNO-Hb(FeII) was converted to GSNO.

Methods

Hbs in PBS (0.5 mM EDTA, pH 7.4) were incubated in air
30 (oxy) or in a tonometer previously evacuated of O₂ (deoxy).

SNO-Hb(FeII)O₂ decomposition was determined by the method of Saville (Saville, B., *Analyst*, 83:670-672 (1958)).

Spontaneous decomposition of SNO-Hb(FeII) was followed spectrophotometrically by formation of Hb(FeII)NO.

- 5 Transnitrosation reactions with glutathione were performed by addition of 100-fold excess glutathione over protein (50 μM), immediate processing of the reaction mixture under anaerobic conditions followed by rapid TCA precipitation, and analysis of RSNO in the supernatant. Rates of NO group
10 transfer were too rapid to measure accurately by the standard methods used in this study.

Example 3: NO-related Interactions with Cysteine Residues of Hb in Physiological Systems

Given that Hb is largely contained within red blood
15 cells, potential mechanisms by which S-nitrosylation of intracellular protein might occur were explored. Incubation of oxygenated rat red blood cells with S-nitrosocysteine resulted in very rapid formation of intracellular SNO-Hb(FeII)O₂ (Figure 3A). Rapid oxidation
20 of Hb was not observed under these conditions.

Intraerythrocytic SNO-Hb also formed when red blood cells were treated with S-nitrosohomocysteine or S-nitrosocysteinyglycine, but not with S-nitrosogluthione (GSNO). Thus, erythrocyte access of RSNOs is thiol group
25 specific. Exposure of oxygenated red blood cells to NO resulted primarily in metHb formation.

Endothelium-derived relaxing factor (EDRF) and hemoglobin

Hb-mediated inhibition of endothelium-dependent relaxations is commonly used as a marker of NO responses.
30 Inasmuch as reactions with either metal or thiol centers of Hb should lead to attenuated NO/EDRF (endothelium-derived relaxing factor) responses, we attempted to elucidate the molecular basis of inhibition. Hb preparations in which

β93 thiol groups had been blocked with N-ethylmaleimide (NEM) or the hemes blocked by cyanmet (FeIIICN)-derivitization were studied in an aortic ring bioassay, and their activities compared with that of native Hb. Both
5 cyanmet-Hb and NEM-Hb caused increases in vessel tone and attenuated acetylcholine (EDRF)-mediated relaxations (Figure 3B). However, native Hb was significantly more effective than either of the modified Hb preparations (Figure 3B). Taken in aggregate, these studies suggest
10 that both the thiol and metal groups of Hb contribute to its NO-related activity. To verify formation of an S-nitrosothiol in Hb, we established a bioassay in which 2 cm segments of thoracic aorta were interposed in Tygon tubing, through which 3 cc of Krebs solution containing Hb (4 μM)
15 and ACh (2 μM) were circulated by roller pump (1.5 cc/min x 5 min). Analysis of the effluent (Gaston, B., et al., (1993)) revealed the formation of SNO-Hb (20 ± 4 nM) in 5 of 5 experiments.

A. S-nitrosylation of intraerythrocytic Hb

20 Incubation of rat erythrocytes with S-nitrosocysteine (equimolar to heme (5 mM); phosphate buffer pH 7.4, 25°C) leads to rapid formation of intracellular SNO-Hb(FeII)O₂. MetHb does not form rapidly. Separation of intracellular
25 RSNOS ~~across~~ ^{across} G-25 columns₃ reveals that only a small percentage exists as low molecular weight S-nitrosothiol (e.g. GSNO) at most time points. By 60 min, 3 of the 4 available SH groups of Hb are S-nitrosylated (note that rat Hb contains 4 reactive SH groups). Inset shows spectra of SNO-Hb isolated from rat erythrocytes and related analyses.
30 Spectrum A is that of SNO-Hb isolated from erythrocytes following G-25 chromatography. Treatment of A with dithionite results in reduction of the S-NO moiety, liberating free NO which is autocaptured by deoxy Hb, forming Hb(FeII)NO (note that dithionite simultaneously

deoxygenates Hb) (spectrum C). This spectrum (C) reveals a stoichiometry of ~3 S-NOs per tetramer. The spectrum of Hb(FeII)NO containing 4 NO's per tetramer is shown for comparison (inset, spectrum B).

5 Methods

At shown intervals, red blood cells were pelleted rapidly by centrifugation, washed three times, lysed in deionized water at 4°C, and the cytosolic fraction subjected to rapid desalting ^{across} ~~across~~ G-25 columns.

10 Intracellular SNO-Hb was measured by the method of Saville (Gaston, B., et al., (1992); Stamler, J.S., et al., *Proc. Natl. Acad. Sci. USA*, 89:444-448 (1992)), and confirmed spectroscopically (inset) as described above.

B. Molecular Basis of EDRF/Hb Interaction.

15 The effects of native Hb on EDRF responses were compared with Hb preparations in which the thiol or heme centers had been blocked by alkylation or cyanmet derivitization, respectively. All preparations of Hb elicited contractions; however, those of native Hb (in
20 which both SH and metal centers are free for interaction) were most pronounced. Likewise, acetylcholine (ACh) mediated relaxations were most effectively inhibited by native Hb. Relaxations were inhibited to lesser degrees by cyanmet Hb (CN-Hb) (in which hemes were blocked from
25 reaction) and NEM-Hb (in which thiol groups were alkylated by N-ethylmaleimide). These data illustrate that both heme and β 93SH groups of Hb contribute to reversal of EDRF responses. Direct measurement of SNO-Hb, formed from EDRF under similar conditions, is described in the text.

30 Methods

Descending rabbit thoracic aorta were cut into 3 mm rings and mounted on stirrups attached to force transducers

(model FT03, Grass Instruments, Quincy, MA) for measurement of isometric tone. The details of this bioassay system have been previously described (Stamler, J.S., et al., *Proc. Natl. Acad. Sci. USA*, 89:444-448 (1992)). Cyanmet Hb
5 was prepared from human HbA according to published protocols (Kilbourn, R.G. et al. *Biochem. Biophys. Res. Comm.*, 199:155-162, (1994)). Alkylation of HbA with N-ethylmaleimide was followed by desalting across G-25 Sephadex to remove excess NEM. Removal of unmodified
10 Hbcys β 93 was achieved by passage through Hg-containing affinity columns. The alkylation of free SH groups was verified using 5,5'-dithio-bis[2-nitrobenzoic acid].

Table 1

ADDITIONS	% INCREASE IN TENSION (†)	% ACh RELAXATION (†)
Hb (1 μ M)	40.8 \pm 2.3 (n=7)	31.9 \pm 6.9 (n=7)
NEM-Hb (1 μ M)	29.4 \pm 1.3 ** (n=7)	60.5 \pm 3.9 * (n=7)
CN-Hb (1 μ M)	12.9 \pm 3.0 ** (n=6)	80.7 \pm 1.0 ** † (n=4)
ACh (1 μ M)		98.3 \pm 0.6 (n=10)
*, P<0.01; **, P<0.001, Compared to Hb; †, P<0.001, Compared to ACh		

Example 4: Transduction of SNO-Hb Vasoactivity

Arterial red blood cells contain two physiologically important forms of hemoglobin: Hb(FeII)O₂ and Hb(FeIII) (Antonini et al. (1971)). Arterial-venous differences in the S-nitrosothiol content of intraerythrocytic Hb suggest that the NO group is released during red cell transit. Such findings raise the possibility of functional consequences, perhaps influenced by the redox state of heme and its occupation by ligand. Intriguingly, SNO-Hb(FeII)O₂ was found to possess modest NO-like activity when tested in a vascular ring bioassay. Specifically, the contraction elicited by SNO-Hb(FeII)O₂ was less than that of native Hb(FeII)O₂, indicating that S-nitrosylation partially reverses the contractile effects of Hb (Figure 4A). By comparison, SNO-Hb(FeIII) was found to be a vasodilator (Figure 4A). Notably, free NO was devoid of relaxant activity in the presence of Hb(FeII)O₂ or Hb(FeIII) (not shown).

Red blood cells contain millimolar concentrations of glutathione. As equilibria among RSNOs are rapidly established through RSNO/thiol exchange (Arnette and Stamler, ^{J.S.}~~J.S.~~ *Arch. Biochem. Biophys.*, 318:279-285 (1995)), the vasoactivity of SNO-Hb was reassessed in the presence of glutathione. Figure 4B illustrates that glutathione potentiated the vasodilator activity of both SNO-Hb(FeII)O₂ and SNO-Hb(FeIII). GSNO formation under these conditions (confirmed chemically and in bioassay experiments) appeared to fully account for this effect. Further kinetic analyses revealed that transnitrosation involving glutathione was more strongly favored in the equilibrium with SNO-Hb(FeIII) than SNO-Hb(FeII)O₂ (Figure 4C). Given the findings of steady-state levels of SNO-Hb in red blood cells (Table 2 and Figure 3A), these results suggest that 1) the equilibrium between naturally occurring RSNOs and

Hb(cys β 93) lies toward SNO-Hb under physiological conditions; 2) that transnitrosation reactions involving SNO-Hb and GSH are likely to occur within red blood cells (indeed, we have verified the presence of low molecular weight RSNOs in erythrocytes loaded with SNO-Hb); and 3) that oxidation of the metal center of Hb will shift the equilibrium toward GSNO, thereby potentially influencing bioactivity.

Additional mechanisms of NO group release from SNO-Hb were sought. Arterial-venous differences in levels of SNO-Hb raised the possibility that S-NO bond stability may be regulated by the changes in Hb conformation accompanying deoxygenation. To test this possibility, we compared the rates of NO group release from SNO-Hb(FeII)O₂ and SNO-Hb(FeIII). Deoxygenation was found to enhance the rate of SNO-Hb decomposition (Figure 2B). These rates were accelerated greatly by glutathione in a reaction yielding GSNO (Figure 2B). Our results illustrate that O₂-metal interactions influence S-NO affinity, and suggest a new allosteric function for Hb.

For SNO-Hb to be of physiological importance it must transduce its NO-related activity accross the erythrocyte membrane. We therefore explored this possibility by incubating erythrocytes containing SNO-Hb in physiologic buffer, and measuring the accumulation of extracellular RSNOs over time. Figure 4D illustrates that red blood cells export low molecular weight (trichloroacetic acid ~~precipitable~~ ^{soluble}) S-nitrosothiols under these conditions.

Importantly, the degree of hemolysis in these experiments was trivial (<0.5%), and correction for lysis did not significantly impact on rates of RSNO release. These results establish that an equilibrium exists between low molecular weight and protein RSNOs within the red cell, and that intracellular location is unlikely to be a limiting

factor in the transduction of such NO-related activity to the vessel wall.

A. Concentration-effect responses of different SNO-Hb preparations.

5 Contractile effects of $\text{Hb(FeII)O}_2(\blacktriangle)$ are shown to be partially reversed by S-nitrosylation ($\text{SNO-Hb[FeII]O}_2(\blacksquare)$; $P = 0.02$ by ANOVA vs Hb(FeII)O_2). Oxidation of the metal center of SNO-Hb ($\text{SNO-Hb[FeIII]}(\bullet)$) converts the protein into a vasodilator ($P < 0.0001$ by ANOVA vs SNO-Hb[FeII]O_2),
10 with potency comparable to that of other S-nitrosoproteins (Stamler, J.S., et al., *Proc. Natl. Acad. Sci. USA*, 89:444-448 (1992)). The contractile properties of Hb(FeIII) are shown for comparison (\square); $n = 6-17$ for each data point.

Methods

15 Details of the vessel ring bioassay have been published (Stamler, J.S., et al., *Proc. Natl. Acad. Sci. USA*, 89:444-448 (1992)). SNO-Hb(FeII) O_2 preparations were synthesized with 10-fold excess S-nitrosocysteine (CYSNO) over Hb(FeII)O_2 protein (2% borate, 0.5 mM EDTA, ~15 min
20 incubation), after which desalting was performed across Sephadex G-25 columns. CYSNO was synthesized in 0.5 N HCl, 0.5 mM EDTA and then neutralized (1:1) in 1 M phosphate buffer containing 0.5 mM EDTA. SNO-Hb(FeIII) preparations followed a similar protocol, but used Hb(FeIII) as starting
25 material. The latter was synthesized by treatment of Hb(FeII)O_2 with excess ferricyanide, followed by desalting ^{across} ~~across~~ G-25 columns. SNO-Hb concentrations were verified spectroscopically and the S-nitrosothiol content was determined by the method of Saville (Stamler, J.S., et al.,
30 *Proc. ^{Na-H} Acad. Sci. USA*, 89:444-448 (1992)). The S-NO/tetramer stoichiometry for both SNO-Hb preparations was ~2. Oxidation of the heme was undetectable by uv-spectrophotometric methods.

B. Potentiation of SNO-Hb effects by Glutathione

Addition of glutathione (100 μ M) to bioassay chambers potentiates the dose-response to both SNO-Hb(FeII)O₂ (■) and SNO-Hb(FeIII) (●) (n=6-12; p<0.0001 for both by ANOVA, 5 compared with the respective tracings in a). Glutathione had a transient affect on baseline tone in some experiments, and did not significantly influence the response to Hb(FeII)O₂ (▲).

C. Transnitrosation between SNO-Hb and Glutathione

10 Rates of NO group transfer from SNO-Hb (100 μ M) to glutathione (10 mM) are displayed for SNO-Hb(FeII)O₂ (oxy) and SNO-Hb(FeIII) (met) (n=5). Data are presented as the amount of GSNO formed relative to the starting SNO-Hb concentration. The transfer is more rapid for SNO- 15 Hb(FeIII) than SNO-Hb(FeII)O₂ (p<0.002 by ANOVA), suggesting that the GSNO/SNO-Hb equilibrium is shifted toward GSNO by formation of met Hb.

Methods

Thiol/SNO-Hb exchange, forming GSNO, was verified 20 chemically (Stamler, J.S., et al., *Proc. Natl Acad. Sci. USA*, 89:444-448 (1992)) following trichloroacetic acid precipitation (n=5). These results were verified in separate experiments by measuring the residual SNO-Hb concentration, following separation of reaction mixtures 25 accross G-25 columns.

D. Export of S-nitrosothiols by red blood cells

Human red blood cells containing SNO-Hb are shown to export low molecular weight RSNOs over time. Hemolysis, which ranged from 0-<0.5% over one hour and did not 30 correlate with rates of RSNO release, could account for only a trivial fraction of the measured extracellular RSNO.

Methods

Packed human red blood cells were obtained by centrifugation, washed, and resuspended in phosphate buffered saline containing 5 mM SNO-CYS (0.5 mM EDTA, pH 7.4) for one hour. This results in a red cell preparation containing SNO-Hb (FeII/FeIII mixture) with a stoichiometry of 0.5 S-NO/tetramer. The red blood cells were then washed repeatedly to remove residual CYSNO (verified), and incubated in Krebs' solution (1:4). The accumulation of extracellular RSNO was measured over time by the method of Saville (Saville, B., *Analyst*, 83:670-672 (1958)). Hemolysis was determined by spectral analysis of red blood cell supernatants following centrifugation.

Example 5: SNO-Hb Bioactivity In Vivo

Systemic administration of cell-free Hb results in hypertensive responses which have been attributed to NO scavenging by the heme (Vogel, W.M., et al., *Am. J. Physiol.*, 251:H413-H420 (1986); Olsen, S.B., et al., *Circulation*, 93:329-332 (1996)). To determine if SNO-Hb is free of this adverse affect, and to explore if *in vitro* mechanisms of NO release extend to the *in vivo* circumstance, we compared responses to Hb and SNO-Hb infused as a bolus into the femoral vein of anesthetized rats. As illustrated in Figure 5, Hb(FeII)O₂ (200 nmol/kg) caused an increase in mean arterial pressure of 20 ± 3 mmHg (n = 4; P < 0.05). In contrast, SNO-Hb(FeII)O₂ did not exhibit hypertensive effects and SNO-Hb(FeIII) elicited hypotensive responses (Figure 5). Thus, the profiles of these compounds *in vivo* closely resemble those seen *in vitro* (Figure 4A). Moreover, to demonstrate that the physiological responses of red cells are comparable to those of cell-free Hb preparations, erythrocytes containing SNO-Hb were injected into the femoral vein of rats

pretreated with L-NMMA (50 mg/kg) to deplete endogenous RSNOs. At levels of SNO-Hb comparable to those found in the normal rat (0.1-0.5 μ M), SNO-Hb containing red blood cells elicited hypotensive responses (8 ± 1 mm Hg; mean \pm SEM; n=9), whereas native (SNO-Hb depleted) red blood cells did not ($P=0.001$). These changes in mean blood pressure of ~ 10% are on the order of those that differentiate normotension from hypertension in man, and in the therapeutic range of some antihypertensive regimens. The effects of both Hb and SNO-Hb -- whether cell-free or contained within red cells -- were transient, suggesting that S-nitrosylation of Hb and metabolism of SNO-Hb may be occurring *in vivo*, with consequent restoration of blood pressure. The bioactivity of SNO-Hb in blood, where S-NO/heme stoichiometries approach 1:50,000, is a dramatic illustration of the resistance of this NO-related activity to Hb(Fe) inactivation.

In vivo effects of cell-free Hb and SNO-Hbs

Administration of 2-200 nmol/kg Hb(FeII)O₂ (as a bolus) into the femoral vein of a Sprague-Dawley rat is shown to increase mean arterial pressure in a dose-dependent manner. At 200 nmol/kg, mean arterial pressure increased by 25 mm Hg (20 ± 3 mmHg; n=4; $P < 0.05$). Elevations in blood pressure reversed within 10-15 min. SNO-Hb(FeII)O₂ infusions (over the same dose range) are shown to ameliorate Hb(FeII)O₂-induced hypertension without causing overt changes in blood pressure. A similar response was seen at higher doses. By comparison, SNO-Hb(FeIII) infusions caused a significant fall in mean arterial pressure (pre 108 ± 4 mm Hg; post 74 ± 6 mm Hg, n = 5; $P < 0.05$) at the highest dose (200 nmol/kg). Hypotensive responses tended to be transient with blood pressure normalizing over 10 minutes. A fall in blood

pressure was also seen with injection of erythrocytes containing SNO-Hb.

Methods

5 Rats were anesthetized by intraperitoneal injection of pentobarbital and the femoral arteries and veins accessed by local cut down. The artery was then cannulated and the blood pressure monitored continuously using a Viggo Spectramed pressure transducer attached to a Gould
10 recorder. An IBM PC (DATA Q Cudas) was used for data acquisition.

Example 6: Loading of Red Blood Cells With S-Nitrosothiols

Incubation of rat erythrocytes with S-nitrosocysteine (equimolar to heme (5mM); phosphate buffer pH 7.4, 25°C)
15 leads to rapid formation of intracellular S-nitrosothiols. MetHb does not form rapidly. Separation of cell content across G-25 columns~~s~~ establishes the formation of
intraerythrocytic low molecular weight S-nitrosothiol e.g. S-nitrosoglutathione, (GSNO). By 2 min., one can achieve
20 as much as millimolar GSNO.

Method for assay of RSNO

S-nitrosocysteine (5 mM) treated red blood cells are pelleted rapidly by centrifugation, washed three times, lysed in deionized water at 4°C, and the cytosolic fraction
25 subjected to rapid desalting across G-25 columns. Intracellular RSNO is measured by the method of Saville and can be confirmed spectroscopically.

Effects on blood pressure from loaded red blood cells

Red blood cells treated with S-nitrosocysteine (to produce SNO-RBCs) and introduced into the femoral vein of a Sprague-Dawley^{rat} decreased mean arterial pressure in a dose-
5 dependent manner. For red blood cells in which SNO-Hb was

assayed at 0.3 μ M (the endogenous in vivo SNO-Hb concentration), arterial pressure decreased by 8 ± 1 mm Hg (mean \pm SEM for 9 experiments; $p < 0.001$ compared to untreated red blood cell controls). For red blood cells in
5 which SNO-Hb was assayed at 0.5 μ M, arterial pressure decreased by 10 mm Hg. For red blood cells in which SNO-Hb was assayed at 0.1 μ M (a sub-endogenous SNO-Hb concentration), arterial pressure decreased by 6 mm Hg. The administration of untreated red blood cells caused no
10 effect or a slight increase in arterial blood pressure. Administration of L-monomethyl-L-arginine (L-NMMA; 50 mg/kg) caused an increase in blood pressure of about 20 mm Hg. Changes in blood pressure from a bolus administration of loaded red blood cells lasted 15-20 minutes.

15 Further methods

Rats were anesthetized by intraperitoneal injection of pentobarbital and the femoral arteries and veins accessed by local cut down. The artery was then cannulated and the blood pressure monitored continuously using a Viggo
20 Spectramed pressure transducer attached to a Gould recorder. An IBM PC (DATA Q Codas) was used for data acquisition.

Example 7: Effects of SNO-Hb on Coronary Vasodilation,
Coronary Flow and Blood Pressure

25 SNO-Hb was synthesized as described in Example 4A. Completion of the reaction was determined as described in Example 4A. Twenty-four healthy mongrel dogs (25-30 kg) were anesthetized with intravenous thiamylal sodium (60-80 mg/kg) and subjected to left thoracotomy in the fourth
30 intercostal space. The left circumflex coronary artery distal to the left atrial appendage was minimally dissected. A pair of 7-MHz piezoelectric crystals (1.5 X 2.5 mm, 15-20 mg) was attached to a Dacron backing and

sutured to the adventitia on opposite surfaces of the dissected vessel segment with 6-0 prolene. Oscilloscope monitoring and on-line sonomicrometry (sonomicrometer 120-2, Triton Technology, San Diego, CA) were used to ensure
5 proper crystal position. A pulse Doppler flow probe (10 MHz, cuff type) was implanted distal to the crystals. An inflatable balloon occluder was also placed distal to the flow probe. All branches of the circumflex artery between the crystals and the occluder were ligated. Heparin
10 sodium-filled polyvinyl catheters were inserted into the left ventricular cavity via the apex, into the left atrium via the atrial appendage, and into the ascending aorta via the left internal thoracic artery. The catheters, tubing, and wires were tunnelled to a subcutaneous pouch at the
15 base of the neck.

After a 10 to 15 day recovery period, the catheters and wires were exteriorized under general anesthesia, and 2-3 days later, each dog was given a bolus injection of SNO-Hb (0.4 mg) to evaluate vascular response. Two dogs
20 that demonstrated <5% dilation of epicardial coronary vessels were excluded from subsequent studies, and two were excluded because of other technical reasons.

Dogs were trained and studied while loosely restrained and lying awake in the lateral recumbent position. The
25 laboratory was kept dimly illuminated and quiet. Aortic pressure, left ventricular end-diastolic pressure dP/dt external coronary diameter and coronary flow were monitored continuously. In 10 dogs, 0.1 ml of SNO-Hb solution, 50 nM/kg, was injected via the left atrial catheter. To
30 verify potential effects of solvent on vasculature, 0.1 ml injections of 30% ethanol in distilled water were given as vehicle control. Between injections, phasic coronary blood flow and coronary artery diameter were allowed to return to preinjection levels (minimum 15 minutes). Allowing a 15
35 minute period between injections resulted in no

a
modification of repeated ^{dose}~~doses~~ injections. To assess the direct and potential flow mediated indirect vasodilation effects of SNO-Hb on the conductance vessels, the dose was repeated in 6 of 10 dogs with partial inflation of the
5 adjustable occluder to maintain coronary blood flow at or slightly below preinjection levels. The response to acetylcholine chloride (Sigma Chemical) was assessed in another group of 10 dogs following a similar protocol to that used for SNO-Hb.

10 Epicardial coronary diameter, coronary blood flow, heart rate, and aortic and left ventricular end-diagnostic pressures were compared before and after each SNO-Hb injection. The maximum changes in coronary dimension and blood flow were expressed as a function of increasing doses
15 of SNO-Hb. The response of coronary dimension to increasing doses followed a characteristic sigmoid dose-response curve that could be described by the following equation

$$Effect = \frac{maximal\ effect\ x\ dose}{K_D + dose}$$

where K_D is the drug-receptor complex dissociation constant
20 and is the dose at which 50% of the maximum response (EC_{50}) is achieved. In each animal, a nonlinear least-squares regression ($r^2 > 0.90$) was performed on the dose-response data. The regression was constrained to the above
equation. From the regression, values for maximum response
25 and K_D were obtained for each individual animal. The mean of these values was then calculated to obtain an average K_D and maximum response for the study group. These values were used to generate a mean curve, which was plotted with the mean dose-response values. (See Figures 6A-6F.)

Example 8: Endogenous Levels of S-nitrosohemoglobin and Nitrosyl(FeII)-Hemoglobin in Blood

To determine if SNO-Hb is naturally occurring in the blood, and if so, its relationship to the O₂ transport capacity and nitrosylated-heme content of red cells, we developed an analytical approach to assay the S-nitrosothiol and nitrosyl-heme content of erythrocytes (Table 2). Arterial blood was obtained from the left ventricle of anesthetized rats by direct puncture and a 10 venous blood was obtained from the ^{jugular} ~~jugular~~ vein and inferior vena cava. Hb was then purified from red cells and assayed for RSN₂O and (FeII)NO content. Arterial blood contained significant levels of SNO-Hb, whereas levels were virtually undetectable in venous blood (Table 2).

15 Measurements made 45 minutes after infusion of the NO synthase inhibitor N^ω-monomethyl-L-arginine (L-NMMA) (50 mg/kg), showed a depletion of SNO-Hb as well as total Hb-NO (82 and 50 ± 18%, respectively; n=3-5; p<0.05). These data establish the endogenous origin of SNO-Hb, although some

20 environmental contribution is not excluded. The arterial-venous distribution seen for SNO-Hb was reversed in the case of Hb(FeII)NO, which was detected in higher concentrations in partially deoxygenated (venous) erythrocytes (Table 2). Accordingly, the proportion of

25 nitrosylated protein thiol and heme appears to depend on the oxygenation state of the blood. Consistent with these findings, Wennmalm and coworkers have shown that Hb(FeII)NO forms mainly in venous (partially deoxygenated) blood (Wennmalm, A., et al., *Br. J. Pharmacol.*, 106(3):507-508

30 (1992)). However, levels of Hb(FeII)NO *in vivo* are typically too low to be detected (by EPR) and SNO-Hb is EPR-silent (*i.e.* it is not paramagnetic). Thus, photolysis-chemiluminescence represents an important technological advance, as it is the first methodology

capable of making quantitative and functional assessments of NO binding to Hb under normal physiological conditions.

Methods

Blood was obtained from the left ventricle (arterial) and jugular vein (venous) of anesthetized Sprague-Dawley rats. Comparable venous values were obtained in blood from the inferior vena cava. Red blood cells were isolated by centrifugation at 800 g, washed three times in phosphate buffered saline at 4°C, lysed by the addition of 4-fold excess volume of deionized water containing 0.5 mM EDTA, and desalted rapidly across G-25 columns according to the method of Penefsky at 4°C. In 24 rats, Hb samples were divided in two aliquots which were then treated or not treated with 10-fold excess HgCl_2 over protein concentration as measured by the method of Bradford. Determinations of SNO-Hb and Hb(FeII)NO were made by photolysis-chemiluminescence as described below. In 12 additional rats, further verification of the presence of SNO-Hb was made by assaying for nitrite after HgCl_2 treatment. Specifically, samples (with and without HgCl_2) were separated across Amicon-3 (Centricon filters, m.w. cut off 3,000) at 4°C for 1 h, and the low molecular weight fractions collected in airtight syringes containing 1 μM glutathione in 0.5 N HCl. Under these conditions, any nitrite present was converted to S-nitrosoglutathione, which was then measured by photolysis-chemiluminescence (detection limit ~ 1 nM). SNO-Hb was present in all arterial samples, and levels determined by this method (286 ± 33 nM) were virtually identical to and not statistically different from those shown in Table 2. In venous blood, SNO-Hb was undetectable (0.00 ± 25 nM); levels were not statistically different from those given above.

Method for assay of S-nitrosohemoglobin

A highly sensitive photolysis-chemiluminescence methodology was employed. A somewhat similar assay has been used for measuring RSNOs (S-nitrosothiols) in biological systems (Gaston, B., et al., (1993); Stamler, J.S., et al., (1992)). The method involves photolytic liberation of NO from the thiol, which is then detected in a chemiluminescence spectrometer by reaction with ozone. The same principle of operation can be used to cleave (and measure) NO from nitrosyl-metal compounds (Antonini, E. Brunori, M. *In Hemoglobin and Myoglobin in Their Reactions with Ligands*, American Elsevier Publishing Co., Inc., New York, pp. 29-31 (1971)). With adjustment of flow rates in the photolysis cell, complete photolysis of the NO ligand of Hb(FeII)NO could be achieved. Standard curves derived from synthetic preparations of SNO-Hb, Hb(FeII)NO, and S-nitrosoglutathione were linear ($R > 0.99$), virtually superimposable, and revealing of sensitivity limits of approximately 1 nM. Two analytical criteria were then found to reliably distinguish SNO-Hb from Hb(FeII)NO: 1) signals from SNO-Hb were eliminated by pretreatment of samples with 10-fold excess HgCl_2 , while Hb(FeII)NO was resistant to mercury challenge; and 2) treatment of SNO-Hb with HgCl_2 produced nitrite (by standard Griess reactions) in quantitative yields, whereas similar treatment of Hb(FeII)NO did not. UV/VIS spectroscopy confirmed that NO remained attached to heme in the presence of excess HgCl_2 .

Table 2

Endogenous levels of S-nitrosohemoglobin and
nitrosyl(FeII)-hemoglobin in blood

	Site	SNO-Hb (nM)	Hb(FeII)NO (nM)
5	Arterial	311 ± 55*	536 ± 99 †
	Venous	32 ± 14	894 ± 126

* P < 0.05 vs venous; † P < 0.05 for paired
samples vs venous

10 Equivalents

Those skilled in the art will know, or be able to
ascertain using no more than routine experimentation, many
equivalents to the specific embodiments of the invention
described herein. These and all other equivalents are
15 intended to be encompassed by the following claims.